



Product Datasheet

Product Name	TIE2 Fc Chimera Human Recombinant
Cata No	CB500854
Source	<i>Insect Cells</i>
Synonyms	Angiopoietin-1 receptor precursor, Tyrosine-protein kinase receptor TIE-2, hTIE2, Tyrosine-protein kinase receptor TEK, p140 TEK, Tunica interna endothelial cell kinase, CD202b, VMCM, VMCM1, TIE2.

Description

TIE-1 (tyrosine kinase with Ig and EGF homology domains 1) and TIE-2/Tek comprise a receptor tyrosine kinase (RTK) subfamily with unique structural characteristics: two immunoglobulin-like domains flanking three epidermal growth factor (EGF)-like domains and followed by three fibronectin type III-like repeats in the extracellular region and a split tyrosine kinase domain in the cytoplasmic region. These receptors are expressed primarily on endothelial and hematopoietic progenitor cells and play critical roles in angiogenesis, vasculogenesis and hematopoiesis. Human TIE-1 cDNA encodes a 1122 amino acid (aa) residue precursor protein with an 18 residue putative signal peptide, a 726 residue extracellular domain and a 353 residue cytoplasmic domain. Two ligands, angiopoietin-1 (Ang1) and angiopoietin-2 (Ang2), which bind TIE-2 with high-affinity have been identified. Ang2 has been reported to act as an antagonist for Ang1. Mice engineered to overexpress Ang2 or to lack Ang1 or Tie-1 display similar angiogenic defects.

Soluble TIE-2 Human Recombinant fused with the Fc part of human IgG₁ produced in baculovirus is a monomeric, glycosylated, polypeptide containing 730 amino acids and having a total molecular mass of 250 kDa. Human TIE-2/Fc monomer has a calculated molecular mass of approximately 125 kDa. As a result of glycosylation, the recombinant

protein migrates as an approximately 140 kDa protein in SDS-PAGE under reducing conditions. The TIE2 Fc Chimera is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Biological Activity

Measured in a functional ELISA assay. When TIE-2/Fc is immobilized at 4 µg/mL (100 µl/well), it binds rh Angiopoietin-2 with a linear range of 2 -100 ng/ml.

Purity

Greater than 90.0% as determined by:
(a) Analysis by RP-HPLC.
(b) Analysis by SDS-PAGE.

Formulation

TIE-2 Fc Chimera was lyophilized from a concentrated (1 mg/ml) sterile solution containing 1x PBS.

Reconstitution

It is recommended to reconstitute the lyophilized TIE-2 Fc Chimera in sterile water not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized sTIE-2 although stable at room temperature for 3 weeks, should be stored

*** For Non-Clinical Research Use Only ***



California Bioscience

83103 Avenue 48, Ste.1B #204
Coachella, CA 92236 USA
Phone : +1.6268339877
Email : info@cali-bio.com

desiccated below -18°C . Upon reconstitution TIE-2 should be stored at 4°C between 2-7 days and for future use below -18°C .

For long term storage it should be stored in the presence of a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.

Product Data Sheet

*** For Non-Clinical Research Use Only ***