



## Product Datasheet

<b>Product Name</b>	B-Cell Maturation Antigen Human Recombinant
<b>Cata No</b>	CB501332
<b>Source</b>	<i>Escherichia Coli.</i>
<b>Synonyms</b>	BCMA, CD269, Tumor Necrosis Factor Receptor Superfamily Member 17, BCM, TNFRSF17, B-cell maturation protein, CD269 antigen.

### Description

TNFRSF17 is a receptor for tnfsf13b/blys/baff and tnfsf13/april. TNFRSF17 promotes b-cell survival and plays a role in the regulation of humoral immunity. TNFRSF17 activates nf-kappa-b and jnk. TNFRSF17 is a member of the TNF-receptor superfamily. TNFRSF17 is expressed in mature B lymphocytes, and is involved in B cell development and autoimmune response. TNFRSF17 specifically binds to the tumor TNFSF13B/TALL-1/BAFF, which causes NF-kappaB and MAPK8/JNK activation. TNFRSF17 binds to a variety of TRAF family members, and therefore transduces signals for cell survival and proliferation. TNFRSF17 is a type III membrane protein having 1 extracellular cysteine rich domain. Within the TNFRSF, it shares the highest homology with TACI. BCMA and TACI have both been shown to bind to APRIL and BAFF, members of the TNF ligand superfamily. BCMA expression has been found in immune organs. TNFRSF17 appears to be localized to the Golgi compartment. The binding of BCMA to APRIL or BAFF has been shown to stimulate IgM production in peripheral blood B cells and increase the survival of cultured B cells.

TNFRSF17 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 50 amino acids and having a molecular mass of 5.3 kDa.

The TNFRSF17 is purified by standard chromatographic techniques.

### Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

### Purity

Greater than 98.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

### Formulation

1mg of TNFRSF17 Human contain 20mM PBS, pH-7.4, and 130mM NaCl.

### Reconstitution

It is recommended to reconstitute the lyophilized TNFRSF17 in sterile 18MΩ-cm H<sub>2</sub>O not less than 100μg/ml, which can then be further diluted to other aqueous solutions.

### Stability

Lyophilized TNFRSF17 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution TNFRSF17 should be stored at 4°C between 2-7 days and for future use below -18°C.

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

**Please prevent freeze-thaw cycles.**

### Sequence

AGQCSQNEYF DSHLHACIPC QLRCSSNTTPP  
LTCQRYCNAS VTNSVKGTNA

**\* For Non-Clinical Research Use Only \***



# California Bioscience

---

83103 Avenue 48, Ste.1B #204  
Coachella, CA 92236 USA  
Phone : +1.6268339877  
Email : [info@cali-bio.com](mailto:info@cali-bio.com)

## Product Datasheet

**\* For Non-Clinical Research Use Only \***